

## Abstracts of free papers presented at the annual meeting of the Obstetric Anaesthetists' Association, Sheffield, 7-8 June, 2007

(The presenter is underlined)

### ORAL PRESENTATIONS

- O01** **Cost effectiveness of using a cell saver in a large obstetric unit** S5  
M King, I Wrench, P Whiting
- O02** **Prophylactic bilateral internal iliac artery catheterisation for control of haemorrhage during elective caesarean section for grade 4 anterior placenta praevia: a series of six cases** S5  
M Hariprasad, S Monaghan, R Sharma, R Wadsworth
- O03** **Determination of the dose-response relationship of spinal bupivacaine, levobupivacaine and ropivacaine, combined with sufentanil, during anaesthesia for caesarean section** S6  
E Roofthoof, R Dreelinck, J Dubois, N Berends, M Van de Velde
- O04** **Carbonated lidocaine v. levobupivacaine for extending epidural analgesia for emergency caesarean section** S6  
J Allam, C Hemingway, S Malhotra, S M Yentis
- O05** **Intrathecal initiation of labour analgesia using fentanyl or diamorphine: is more always better?** S7  
S Dinesh, N Sabir, A Majumder, P N Robinson, D J A Vaughan
- O06** **The hypotensive and fetal heart rate response to low-dose epidurals: analysis of a randomised trial dataset** S7  
K Duhig, C MacArthur, A May, A H Shennan
- O07** **Combined spinal epidural analgesia during labour: is a prophylactic intravenous crystalloid fluid load required to prevent hypotension?** S8  
E Roofthoof, Y Van Dormael, E Vandermeersch, R Devlieger, M Van de Velde
- O08** **Maternal cardiac output changes after crystalloid or colloid cohydration following spinal anaesthesia for elective caesarean section** S8  
S McDonald, R Fernando, K Ashpole, M Columb
- O09** **Effects of crystalloid preload and spinal anaesthesia for caesarean section on maternal haemodynamics using LiDCOplus system** S9  
N Daniel, P Barclay, R Sieunarine, A Carlin, Z Alfirevic
- O10** **Hypotension following low dose combined spinal-epidural anaesthesia for caesarean section: left lateral versus supine tilted position** S9  
O Adekanye, R Sivasankar, R E Collis
- O11** **Effect of intravenous phenylephrine or ephedrine on the ED50 of intrathecal bupivacaine with fentanyl for caesarean section** S10  
C Hennebry, G M Stocks, P Belavadi, K Mordani, J Barnes, M Columb, G Lyons
- O12** **Acid-base balance in an obstetric population: establishing a normal range** S10  
D Thorp-Jones, S Yadhore, R E Collis
- O13** **Does eating during labour influence obstetric outcome?** S11  
B Liu, G O'Sullivan, D Hart, M Waterstone, A H Shennan
- O14** **Cardiac maternal deaths: missing data in the CEMACH/CEMD reports over 30 years** S11  
S Malhotra, S M Yentis
- O15** **A blinded study of the sensitivity of epidural catheter aspiration in identifying intrathecal epidural catheter placement** S12  
C R W Baird, S R McKechnie
- O16** **Stability of adrenaline in pH-adjusted lidocaine** S12  
J Allam, H Gill, C Tuleu, P Knowlden, E Smith, S M Yentis
- O17** **Obstetric workload of non-obstetric consultant anaesthetists: a national OAA-approved survey** S13  
S Sagadaj, S Maguire
- O18** **Publication of papers arising from abstracts presented at OAA annual meetings 1994-2004** S13  
R Srivastava, K Rahman, S M Yentis

### POSTER PRESENTATIONS

- P01** **Multimodal analgesia for caesarean section does not need intrathecal diamorphine** S14  
V Sood, M Porter, S Chaudhari, A Johal
- P02** **How painful is epidural insertion? What mothers think** S14  
K Ray, M Purva, Z Rafique, I F Russell

<b>P03</b>	<b>An audit of maternal nausea and vomiting following a change from phenylephrine boluses to an infusion</b> <u>M McCormick</u> , A Macfarlane, E McGrady	S15
<b>P04</b>	<b>Temperature loss during elective caesarean section under spinal anaesthesia</b> <u>J Limb</u> , K Adams, B Heidemann	S15
<b>P05</b>	<b>Chemical burns and pressure sores in obstetric anaesthesia: a retrospective audit</b> <u>M Chakravarti-Chattopadhyay</u> , N Patel, R Bell	S16
<b>P06</b>	<b>Complications associated with combined spinal-epidural vs spinal anaesthesia for caesarean section</b> <u>A Allnatt</u> , R Sashidharan	S16
<b>P07</b>	<b>Complications of regional techniques for obstetric analgesia and anaesthesia</b> <u>Worms R</u> , Gill F, Sharpe P	S17
<b>P08</b>	<b>Increasing dural tap rate: is this a national trend?</b> <u>S Gupta</u> , R E Collis, S E Harries	S17
<b>P09</b>	<b>Past history of PDPH and spinal anaesthesia</b> S Gowrie-Mohan, <u>G Hallward</u> , V Luoma, M Cassim	S18
<b>P10</b>	<b>Subarachnoid haemorrhage following dural puncture: case report</b> M Girgis, <u>H Swales</u> , S Hughes, J Hell	S18
<b>P11</b>	<b>Paraparesis following spinal anaesthesia for elective caesarean section: a case report</b> <u>M Rwemi</u> , R K Aggarwal, A Swami	S19
<b>P12</b>	<b>Monitoring of mother and fetus before and during regional analgesia: comparison of three different UK teaching hospitals</b> <u>A Thangamuthu</u> , M Purva	S19
<b>P13</b>	<b>Fetal ST analysis in labour: implications for anaesthetic management</b> <u>R Hamilton</u> , A May, L Powls, J Reid	S20
<b>P14</b>	<b>Fetal heart rate monitoring following spinal anaesthesia is important for anaesthetic management before caesarean section</b> <u>K Suzuki</u> , S Irikoma, S Kokubo	S20
<b>P15</b>	<b>Audit of intrauterine fetal resuscitation before caesarean section for fetal compromise</b> <u>N K Weale</u> , S M Kinsella	S21
<b>P16</b>	<b>Prospective randomised study of external cephalic version for breech presentation at term in nulliparous women: spinal analgesia versus no analgesia</b> C F Weiniger, Y Ginosar, U Elchalal, S Einav, M Nucrietin, P Guage, Y Ezra	S21
<b>P17</b>	<b>Four-year prospective audits (2003-2006) of general anaesthesia and conversion of regional to general anaesthesia for caesarean section</b> <u>M A Rafi</u> , D McIntosh, Z Arfeen, U Misra	S22
<b>P18</b>	<b>Types of anaesthesia for grade 1 and 2 caesarean sections in 2006</b> <u>M A Rafi</u> , Z Arfeen, U Misra	S22
<b>P19</b>	<b>Emergency caesarean section under general anaesthesia: can it be avoided?</b> <u>N G Davies</u> , R E Collis	S23
<b>P20</b>	<b>Fasting in elective caesarean section</b> <u>I Ahmed</u> , H Brooks, N Hickman	S23
<b>P21</b>	<b>General anaesthesia for caesarean section: are intubation problems on the increase?</b> <u>E Plunkett</u> , E Wong	S24
<b>P22</b>	<b>Failed intubation: the tip of the iceberg</b> <u>C Burnett</u> , R Hamilton, J Reid	S24
<b>P23</b>	<b>Spinal, single-shot epidural or general anaesthesia for caesarean delivery</b> <u>A Hasani</u> , B Ay, S Azizi	S25
<b>P24</b>	<b>Effects of gender and pregnancy on ED50 for motor block with intrathecal bupivacaine</b> <u>M Camorcia</u> , G Capogna, M Columb	S25
<b>P25</b>	<b>Training declines in obstetric general anaesthesia</b> <u>R Searle</u> , G Lyons	S26
<b>P26</b>	<b>A comparison of different solutions for volume preloading: prevention of hypotension during epidural anaesthesia for caesarean section</b> <u>E Shifman</u> , I Got	S26
<b>P27</b>	<b>A survey of pre-loading and vasopressor use during regional anaesthesia for caesarean section</b> <u>A McGlennan</u> , N Patel, B Sujith, R Bell	S27
<b>P28</b>	<b>Caesarean section for placenta praevia: are we practising evidence-based medicine?</b> <u>A M McEwen</u> , L Campbell	S27
<b>P29</b>	<b>Haemodynamic changes associated with spinal anaesthesia for caesarean section in severe preeclampsia</b> <u>R A Dyer</u> , AR Reed, JL Piercy, L Schoeman, M F James	S28
<b>P30</b>	<b>Successful use of recombinant factor VIIa in the treatment of life threatening postpartum haemorrhage in a Jehovah's Witness patient</b> <u>W Wight</u> , R Khoju	S28
<b>P31</b>	<b>Prophylactic uterine artery balloon occlusion and embolisation in placenta percreta</b> <u>A Sinha</u> , H Swales, S Hughes, N Hacking	S29

<b>P32</b>	<b>Massive obstetric haemorrhage: moving upstream into the aorta</b> <u>S Humble</u> , P Agustsson, P Clark, S Chakraverty, I Zealley, C Connolly	S29
<b>P33</b>	<b>Survey of cell-salvage use in obstetrics in the UK</b> <u>M Teig</u> , M Harkness, S Catling, V Clark	S30
<b>P34</b>	<b>Triggers for transfusion for vaginal and caesarean deliveries</b> R Akthar, <u>S Sivasubramaniam</u>	S30
<b>P35</b>	<b>Changing patterns in obstetric haemorrhage: experience of a London DGH</b> <u>D N Lucas</u> , P N Robinson	S31
<b>P36</b>	<b>Ethnicity and post-partum haemorrhage</b> <u>L Edger</u> , P N Robinson, K Rao, T Rudra, B A Loughnan	S31
<b>P37</b>	<b>Reducing risk in an ethnic minority: a retrospective observational study</b> <u>K Ray</u> , G Lyons	S32
<b>P38</b>	<b>Are we being called? An audit of anaesthetic referral for pre-assessment of high-risk mothers</b> <u>S Hebard</u> , J A Thurlow	S32
<b>P39</b>	<b>Review of analgesia and pain scores in substance-misusing women requiring caesarean section</b> <u>M Woolnough</u> , I Wrench	S33
<b>P40</b>	<b>Management and outcome of parturients with cardiac disease: a one year retrospective survey</b> <u>S Al-Rawi</u> , P Gill	S33
<b>P41</b>	<b>Is there a role for enoximone in the management of pregnant women with dilated cardiomyopathy?</b> <u>E Hossenbaccus</u> , C Robinson, R Wadsworth, A Vohra	S34
<b>P42</b>	<b>Use of vasopressin during caesarean section in parturients with idiopathic pulmonary artery hypertension</b> <u>L Price</u> , V Sodhi, M Lucey, D Adamson, C Nelson-Piercy, P Forrest, L S Howard	S34
<b>P43</b>	<b>The use of oxytocin in parturients with cardiac disease: a case series</b> <u>S Wray</u> , V Sodhi, D Adamson, M K Dhanjal, C Nelson-Piercy	S35
<b>P44</b>	<b>Assessment of pneumonia severity in pregnancy</b> <u>A Macfarlane</u> , E McGrady	S35
<b>P45</b>	<b>Uterotonics and pulmonary oedema: is there a link?</b> <u>L Rees</u> , R E Collis, S Harries	S36
<b>P46</b>	<b>Arnold-Chiari malformation and childbirth</b> <u>G Sridhar</u> , J Evans	S36
<b>P47</b>	<b>National survey of anaesthetic assessment of pregnant women with increased body mass index</b> H Smith, <u>B Varghese</u> , P Gill, H Swales	S37
<b>P48</b>	<b>Critically ill obstetric patients treated in an obstetric intensive care unit</b> E Gollo, <u>S Calva</u>	S37
<b>P49</b>	<b>CSE in morbidly obese parturients</b> <u>A Gobindram</u> , S Gowrie-Mohan, S McNeillis, P U Ramanayake	S38
<b>P50</b>	<b>Preparation of midwives for high risk care: viewpoint of the midwife</b> <u>S Rathinam</u> , A Khan, M Rupasinghe	S38
<b>P51</b>	<b>Patient-controlled epidural analgesia for labour: is anaesthetic and midwifery workload reduced?</b> <u>D Cochran</u> , P Moore	S39
<b>P52</b>	<b>The use of obstetric patient-controlled epidural analgesia: a survey of labour wards across the UK</b> <u>O Boswell</u> , A Cowan, S Pilkington, I Taylor	S39
<b>P53</b>	<b>Discontinuing epidurals in the late stages of labour: a survey among midwives in the North West of UK</b> <u>S Sagadaj</u> , H Vallance, S Maguire	S40
<b>P54</b>	<b>Epidural top-ups for second stage of labour: a survey of midwifery practice in our unit</b> <u>S Rathinam</u> , P Tilakaratna, H Bojahr	S40
<b>P55</b>	<b>Indices of deprivation, perceived pain and epidural uptake in primiparous women undergoing induction of labour</b> <u>J Dolan</u> , J Hislop, P Roche, S Young	S41
<b>P56</b>	<b>Does social deprivation affect epidural requests?</b> <u>A Macfarlane</u> , S Young	S41
<b>P57</b>	<b>Audit of pain relief in labour: how informed are pregnant women of ethnic minority origin attending the maternity unit of a district general hospital?</b> <u>H Goddard</u> , A Obideyi, T Isitt	S42
<b>P58</b>	<b>Service review of intrathecal anaesthesia for caesarean section following epidural analgesia in labour</b> <u>G Thomas-Kattappurathu</u> , I Wrench	S42
<b>P59</b>	<b>Adequacy of anaesthesia for caesarean section after unsuccessful instrumental vaginal delivery</b> <u>A Kathirgamanathan</u> , W J D Whiteley, D G Bogod, D M Levy	S43
<b>P60</b>	<b>Audit of remifentanyl patient-controlled analgesia for labour</b> <u>R Laird</u> , D Hughes, D Hill	S43
<b>P61</b>	<b>Prospective audit of the impact of a new obstetric anaesthesia chart on record keeping</b> <u>C A Slorach</u> , S Neal	S44
<b>P62</b>	<b>Comparison of obstetric anaesthesia study populations with a tertiary referral obstetric population</b> A Macfarlane, <u>H McKay</u> , S Young	S44

<b>P63</b>	<b>Obstetric high dependency facilities: a survey of current practice</b> K Srinivas, S Yadthore, R E Collis	S45
<b>P64</b>	<b>Ramadan: a survey of staff attitudes to fasting during pregnancy</b> Z Apple, S Rathinam, M Bartimeus, H Bojahr, M Razzaque	S45
<b>P65</b>	<b>Obstetric epidurals: not just inserted at night!</b> R Marr, M Millar	S46
<b>P66</b>	<b>Maternal expectations and experiences of labour analgesia: an observational study</b> N Curran, M Holmes, J Holding, M Ryten, L Bromley, B Brandner, R Bell	S46
<b>P67</b>	<b>Maternal satisfaction and knowledge about epidural analgesia: a phone call survey</b> G Vasconcelos, C Oliveira, L Marinho	S47
<b>P68</b>	<b>What do women want? An audit of antenatal information on analgesia for labour</b> G Allan, N Parry, V Skelton	S47
<b>P69</b>	<b>Why obstetric epidural blocks occasionally fail: results from the Australian epidurogram study</b> C Collier	S48
<b>P70</b>	<b>Fruits of our labour: a novel epidural simulator</b> D Raj, R Williamson, D Young, D Russell	S48
<b>P71</b>	<b>Evaluation of three methods of determining the L3-4 lumbar interspace using MR imaging</b> S Yadthore, K Srinivas, G Griffiths, R E Collis	S49
<b>P72</b>	<b>Graphical presentation of labour analgesia</b> T Fernandez, M Mackenzie, J Durbridge, S M Yentis	S49
<b>P73</b>	<b>When to start a caesarean section after an epidural top-up: touch to T5, cold to T4, or something else?</b> N A Richards, J Allam, S Malhotra and S M Yentis	S50
<b>P74</b>	<b>Obstetric continuous spinal analgesia/anaesthesia: a survey of UK practice</b> S Whittaker, M Fletcher, M Wee	S50
<b>P75</b>	<b>Spinal catheters in obstetric anaesthesia: experience from two tertiary centres</b> H Wilson, M Scrutton, M Kinsella, A Pinder, Dresner M	S51
<b>P76</b>	<b>Spinal needle, stylet and Euler buckling</b> C Macartney, D A Hill, A Gallagher, D Carson	S51
<b>P77</b>	<b>Ultrasound-supported epidural catheterisation for emergency labour analgesia</b> S Baglioni, L D'Andrea, A Moretto, M Negri, G Barontini, I Buquicchio, R Fumagalli, A Pesenti	S52
<b>P78</b>	<b>Variation in the relative position of the conus medullaris in the neutral and lateral decubitus flexed positions of the lumbar spine in healthy females: a magnetic resonance imaging study</b> U Misra, P Evans, T Hildreth, S England	S52
<b>P79</b>	<b>Desperately seeking caesarean: search terms used by anaesthetists when searching electronic databases</b> R Srivastava, S M Yentis	S53
<b>P80</b>	<b>Synchronisation of maternity unit wall clocks</b> M A Rafi, U Misra	S53
<b>P81</b>	<b>Audit of clocks, watches and timepieces on the delivery suite</b> A Combeer	S54
<b>P82</b>	<b>Levels of care in obstetrics (LOCOBS): level the playing (or paying) field</b> S Wheatly, V McGill, M Columb, W De Mello	S54