

1-1 Obstetric Cardiac Arrest v.2

Alterations in maternal physiology and exacerbations of pregnancy related pathologies must be considered. Priorities include calling the appropriate team members, relieving aortocaval compression, effective cardiopulmonary resuscitation (CPR), consideration of causes and performing a timely emergency hysterotomy if ≥ 20 weeks

START

- 1 Confirm cardiac arrest -and- call for help. Declare 'Obstetric cardiac arrest'**
 - ▶ Team for mother (at any gestation) and team for neonate if ≥ 22 weeks
- 2 Lie flat, apply manual uterine displacement to the left if ≥ 20 weeks or uterus palpable at or above umbilicus**
 - ▶ Or left lateral tilt (from head to toe at an angle of 15-30° on a firm surface)
- 3 Start CPR -and- call for cardiac arrest trolley**
 - ▶ Check for reversible causes (**Box A**)
- 4 Identify team leader, allocate roles including scribe**
 - ▶ Note time
- 5 Apply defibrillation pads and check cardiac rhythm** (defibrillation is safe in pregnancy)
 - ▶ If VF / pulseless VT → defibrillation -and- give first adrenaline and amiodarone after 3rd shock
 - ▶ If PEA / asystole → resume CPR -and- give first adrenaline immediately
 - ▶ Check rhythm and pulse every 2 minutes
 - ▶ Repeat adrenaline every 3-5 minutes
- 6 Maintain airway and ventilation**
 - ▶ Give 100% oxygen using bag-valve-mask device
 - ▶ Insert supraglottic airway with drainage port -or- tracheal tube if trained to do so (Intubation may be difficult and airway pressures may be higher)
 - ▶ Apply waveform capnography (ETCO₂) monitoring to airway
 - ▶ If no expired CO₂ → presume oesophageal intubation
- 7 Circulation**
 - ▶ IV access above the diaphragm, if fails or impossible use upper limb intraosseous (IO)
 - ▶ See (**Box B**) for reminder about drugs
 - ▶ Consider extracorporeal CPR (ECPR) if available
- 8 Emergency hysterotomy (perimortem caesarean section)**
 - ▶ Perform by 5 minutes if no return of spontaneous circulation and ≥ 20 weeks gestation, to improve maternal outcome
 - ▶ Perform immediately if maternal fatal injuries or prolonged pre-hospital arrest

Box A: Reversible causes 4Hs and 4Ts (specific to obstetrics)

Hypoxia	Respiratory – Pulmonary embolism (PE) Failed intubation, aspiration Heart failure Anaphylaxis Eclampsia / PET – pulmonary oedema, seizures
Hypovolaemia	Haemorrhage – obstetric (remember concealed), abnormal placentation, uterine rupture, atony, splenic artery/hepatic rupture, aneurysm rupture Distributive – sepsis, high regional block, anaphylaxis
Hypo/hyperkalaemia	Also check blood sugar, sodium, calcium and magnesium levels
Hypothermia	
Tamponade	Aortic dissection, peripartum cardiomyopathy, trauma
Thrombosis	Amniotic fluid embolism, PE, myocardial infarction, air embolism
Toxins	Local anaesthetic, magnesium, illicit drugs
Tension pneumothorax	Risks include trauma, positive pressure ventilation (including general anaesthesia) Can be exacerbated by Entonox / nitrous oxide

Box B: IV drugs for use during cardiac arrest

Fluids	500 ml IV crystalloid bolus
Adrenaline	1 mg IV every 3-5 minutes in non-shockable or after 3 rd shock
Amiodarone	300 mg IV after 3 rd shock
Atropine	0.5 – 1 mg IV up to 3 mg if vagal tone likely cause
Calcium chloride	10% 10 ml IV for Mg overdose, low calcium or hyperkalaemia
Thrombolysis / PCI	For suspected massive pulmonary embolism / MI
Tranexamic acid	1g if haemorrhage suspected
Intralipid	1.5 ml/kg IV bolus and 15 ml/kg/hr IV infusion